

CUSTOMIZE



GO WITH THE VEIN FLOW

AND INJECT TOWARD THE HEART
Always inject into a vein and never into the skin.
Veins are close to the surface of the skin and spring to the touch.
Arteries are located deeper and are not as visible as veins.
If you hit an artery, it will bleed profusely. Blood will not clot. Blood will not clot. Blood will not clot.
If this happens, apply direct pressure and medical attention if the bleeding does not stop.

Arms usually have three accessible veins.
To find a vein, try using a soft tie to constrict the arm.
Use a safe tie so that you can easily pull the tie off after injecting.
If possible, release the tie after your injection (though, not before you push in your tie).
If you are afraid you will injure yourself, ask a friend to help you push your tie in, push your tie in, push your tie in.
Add your SSP for loan requests and other supplies.

TRY BOTH ARMS
BEFORE GOING TO THE WHISKY, BEEF, AND NECK.

Beetle Up
the MALXONE
www.bvfyg.com
QR Code
© 2014 Beetle Up
www.bvfyg.com

BEVEL DOWN

THE BEVEL IS THE SHARPER EDGE AT THE POINT OF THE SYRINGE
The only time the bevel should face down is when drawing up from a cooker.
You can protect the bevel by using a cotton filter.
100% cotton filter preferred.

BEVEL UP
Position the syringe with the bevel facing up so the sharp tip can pierce the skin first, easing the way for the rest of the needle.

Beetle Up
the MALXONE
www.bvfyg.com
QR Code
© 2014 Beetle Up
www.bvfyg.com

CHASING WITH FOIL

ROCK/CRACK METH
Prepare a foil rectangle that is 1.5 inches wide and 2 inches long (width of your hand).
If the foil has a shiny, plastic coating, pre-burn the foil by holding the shiny side over a flame.
With the dull side facing down, place drugs on the foil and heat from the bottom with a lighter or heat gun. The lighter or heat gun should be a few inches below the foil.

CHASING WITH FOIL
Once the substance has liquefied, use a glass straw or a straw made of foil to chase the smoke.
Take it slow, especially if it's your first time smoking or you haven't smoked in a while.
Remember that it's all possible that it's all possible when smoking, so give yourself a few minutes to see how you feel after that first inhale.

Beetle Up
the MALXONE
www.bvfyg.com
QR Code
© 2014 Beetle Up
www.bvfyg.com

GLASS STRAIGHT STEM

CRACK, ROCK COCAINE
Bress screens are best. Add for them at your SSP.
If you are using a scribbler for a screen, cocaine can break apart and burn your mouth and throat when it gets hot.
Pre-burn your scribbler to release toxic fumes.
If sharing a pipe, use your own mouthpiece and a new filtering cover or rubber band(s).

GLASS BUBBLE PIPE
METH, FENTANYL
Inhale slow and exhale fast.
Heat the pipe from the bottom, but it will damage your hands.
Heat the pipe from side/rear and keep the flame moving around the bowl so you melt the drags instead of the pipe. This prevents the pipe from cracking and wastes less drugs.

Beetle Up
the MALXONE
www.bvfyg.com
QR Code
© 2014 Beetle Up
www.bvfyg.com

BOOTY BUMPING

Mix drugs with 1-2 ml of clean water.
Use a syringe to draw up the mixture.
Inject into the buttock muscle.
Break the point off a syringe (dispose of it safely) or use a lube applicator.
Lube applications are more effective because they are designed to deliver fluid to the rectal cavity. It also holds more liquid than a syringe.

WATER DRIPPING
Water dripping is more effective than sniffing (liquid helps absorption).
Crush drugs as much as possible to reduce irritation.
Water from best to worst: saline, sterile, distilled, bottle or tap.
For a sweeter solution, add more water.
Use a syringe to draw up the mixture.
Break the point off a syringe (dispose of it safely) or use a dropper.
P.S. This only works for drugs that dissolve in water. Water dripping does not work for crack or pills.

Beetle Up
the MALXONE
www.bvfyg.com
QR Code
© 2014 Beetle Up
www.bvfyg.com

IF DRUGS ARE IN YOUR ORBIT, GO WITH MALXONE FENTANYL & NALOXONE

The drug supply is increasing and you need a way of using it that is safe.

If you use sodium, you do your best.

If you are using a syringe, you do your best.

Pre-burn with a lighter, but you can still OD.

Worry, stepping and longer to let in, but you can still OD.

MALXONE comes in a 100 mg/0.1 mg/mL concentration. It's a 100 mg/0.1 mg/mL concentration. It's a 100 mg/0.1 mg/mL concentration. It's a 100 mg/0.1 mg/mL concentration.

MALXONE comes in a 100 mg/0.1 mg/mL concentration. It's a 100 mg/0.1 mg/mL concentration. It's a 100 mg/0.1 mg/mL concentration. It's a 100 mg/0.1 mg/mL concentration.

There is no such thing as a "low cost/low volume" overdose. Naloxone and Narcan will work on all opioid-reversed overdoses, including fentanyl, carfentanyl, and fentanyl analogs.

Beetle Up
the MALXONE
www.bvfyg.com
QR Code
© 2014 Beetle Up
www.bvfyg.com